

# Asian Resonance

## Kinetics and Mechanism of Oxidation of Some Unsaturated Acids by Quinolinium Fluorochromate in Aqueous Acetic Acid Medium-A Comparative Study

### Abstract

The kinetics of oxidation of unsaturated acids viz. maleic acid and acrylic acid by quinolinium fluorochromate (QFC) have been investigated in acetic acid-water mixture (50% v/v) in presence of perchloric acid. The reaction is of first order each in [oxidant], [substrate] and [acid] for both substrates. Increase in dielectric constant of the medium decreases the rate, while variation in ionic strength has no significant effect on the rate. The reaction does not induce polymerisation of acrylonitrile. The reaction rate have been determined at different temperature and the activation and thermodynamic parameters have been computed. Mechanism consistent with the observed results has been discussed.

**Keywords:** Oxidation, Kinetics, Mechanism, Quinolinium Fluorochromate (QFC), Unsaturated Acids, Epoxide.

### J.V.Singh

Associate Professor  
Deptt. of Chemistry,  
Nehru College,  
Chhibramau, Kannauj (U.P.)  
India

### Introduction

The development of new Cr(VI) reagents for the effective and selective oxidation of organic substrates under mild condition has been going on because Cr(VI) is a versatile oxidant for the oxidation of many types of organic substrates and also due to advancement in the synthesis of complex organic molecules under different reaction conditions<sup>1-4</sup>. Quinolinium fluorochromate (QFC) adds to the select list of newer Cr(VI) reagents as the most significant oxidant for the effective and selective oxidation of organic substrates under mild conditions.<sup>5</sup> There seems to be no reports on the kinetic and mechanistic aspects of oxidation of unsaturated acids viz. maleic acid and acrylic acid by QFC.

### Aim of the Study

Hence the present investigation is undertaken and the kinetic results of oxidation of both substrates with QFC in acetic acid water (1:1 v/v) solvent are reported in this paper.

### Sarvesh Kumar

Assistant Professor,  
Deptt. of Chemistry,  
Nehru College,  
Chhibramau, Kannauj (U.P.)  
India

### Experimental

#### Materials

Quinolinium fluorochromate (QFC) was safely prepared by careful addition of quinoline (Loba-chemie) to a solution of chromium trioxide in 40% HF acid followed by the filtration of yellow orange crystal.<sup>5</sup> Solutions of maleic acid (Loba) and acrylic acid (Loba) were always freshly prepared. The ionic strength was kept constant using a concentrated solution of sodium perchlorate (B.D.H.). Perchloric acid (E. Merck) and all other chemicals (A.R., B.D.H.) were used without further purification. Doubly distilled water was used for all kinetic run.

#### Kinetic Measurements

All the kinetic measurements were carried out in 50% acetic acid-50% water mixture containing HClO<sub>4</sub> and NaClO<sub>4</sub> under pseudo first order conditions by keeping excess of substrate over quinolinium fluorochromate at constant temperature ( $\pm 0.1$ K). The progress of the reaction was followed by estimating unreacted oxidant iodometrically after quenching the reaction. The pseudo-first order rate constant  $k_{obs}$ , were determined from the linear plot of log [oxidant] versus time. Replicate runs showed that the rate constant were reproducible to within  $\pm 3\%$  error.

#### Stoichiometry and Product Analysis


Stoichiometry of the reaction for both the substrates was found to be 1 : 1 as determined by carrying out several sets of experiments with varying amounts of [QFC] largely in excess over [substrate]. The reaction

### Ashish Tomar

Assistant Professor,  
Deptt. of Chemistry,  
Meerut College,  
Meerut (U.P.)  
India

E: ISSN No. 2349-9443

mixture was kept 24 hr to ensure the completion of the reaction. The unreacted quinolinium fluorochromate was estimated iodometrically.



Where, R=COOH for maleic acid and R=H for acrylic acid.

Product analysis under kinetic conditions gave only the epoxide in each case of study, which was identified by periodate test<sup>6</sup>.

### Results and Discussion

In each case, the oxidation followed a first order kinetics both with respect to [substrate] and [oxidant]. Plots of log [QFC] versus time were linear upto at least 80% completion of the reaction. The

# Asian Resonance

pseudo-first order rate constants,  $k_{\text{obs}}$ , do not depend on initial concentration of QFC (Table 1.), establishing first order dependence of the rate on [QFC] in both the cases. At constant [QFC] the rate increases steadily with increase in [substrates] (Table 1) and the plots of log  $k_{\text{obs}}$  versus log [substrate] are linear with unit slope, (Fig. 1) showing first order dependence of the rate on [substrate]. The second order rate constants  $k_2 = k_1[\text{substrate}]$  give the concordant values. The plots of  $1/k_1$  against  $1/[\text{substrate}]$  give a straight line passing through origin (Fig. 2). This confirms first order kinetics with respect to substrate and suggests that no intermediate complex is formed between substrate and oxidant. Further UV-visible spectral studies did not show any evidence for the formation of QFC-substrate complex (Fig. 3).


**Table 1: Dependence of rate on [reactants] at 303 K**

Solvent : AcOH-H<sub>2</sub>O (50-50% v/v)

[NaClO<sub>4</sub>] = 2.0 × 10<sup>-1</sup> mol dm<sup>-3</sup>


[QFC] × 10 <sup>3</sup> (mol dm <sup>-3</sup> )	[Maleic acid] × 10 <sup>2</sup> (mol dm <sup>-3</sup> )	[Acrylic acid] × 10 <sup>2</sup> (mol dm <sup>-3</sup> )	[HClO <sub>4</sub> ] (mol dm <sup>-3</sup> )	Maleic acid k <sub>1</sub> × 10 <sup>5</sup> (s <sup>-1</sup> )	Acrylic acid k <sub>1</sub> × 10 <sup>5</sup> (s <sup>-1</sup> )
2.0	2.0	2.0	0.92	9.20	4.12
3.2	2.0	2.0	0.92	9.08	4.00
4.0	2.0	2.0	0.92	9.24	4.10
6.0	2.0	2.0	0.92	9.16	4.14
6.4	2.0	2.0	0.92	9.28	4.08
8.0	2.0	2.0	0.92	9.12	4.06
2.0	1.0	1.0	0.92	4.64	2.00
2.0	1.6	1.6	0.92	7.36	3.22
2.0	2.0	2.0	0.92	9.20	4.12
2.0	3.0	3.0	0.92	13.60	6.20
2.0	3.2	3.2	0.92	14.70	6.40
2.0	4.0	4.0	0.92	18.30	8.30
2.0	5.0	5.0	0.92	23.00	10.30
2.0	2.0	2.0	0.23	2.30	1.05
2.0	2.0	2.0	0.46	4.62	2.04
2.0	2.0	2.0	0.92	9.20	4.12
2.0	2.0	2.0	1.38	13.86	6.20
2.0	2.0	2.0	1.84	18.44	8.14
2.0	2.0	2.0	2.30	23.08	10.48

**Fig. 1. Plot of  $\log k_1$  vs  $\log$  [unsaturated acid]**  
 [QFC] =  $2.0 \times 10^{-3}$  mol dm $^{-3}$ ; [HClO $_4$ ] = 0.92 mol dm $^{-3}$ ; [NaClO $_4$ ] =  $2.0 \times 10^{-1}$  mol dm $^{-3}$ ; Solvent : AcOH-H $_2$ O (50-50% v/v); Temperature : 303 K.




**Fig. 2. Plot of  $\log k_1^{-1}$  vs [substrate] $^{-1}$**

[QFC] =  $2.0 \times 10^{-3}$  mol dm $^{-3}$ ; [HClO $_4$ ] = 0.92 mol dm $^{-3}$ ; [NaClO $_4$ ] =  $2.0 \times 10^{-1}$  mol dm $^{-3}$ ; Solvent : AcOH-H $_2$ O (50-50% v/v); Temperature : 303 K




**Fig. 3. Electronic spectra of QFC in aqueous acetic acid (50% v/v) (A) without substrate (B) with maleic acid (C) with acrylic acid**



The direct proportionality between  $k_1$  values and  $[\text{HClO}_4]$  indicate first order kinetics with respect to  $[\text{H}^+]$  (Table 1). The plots of  $\log k_1$  versus  $\log [\text{H}^+]$  gave straight line with a slope of unity (Fig. 4).

The linear increase in the rate with acidity suggests the involvement of protonated Cr(VI) species in the rate determining step. The participation of protonated chromium species in Cr(VI) oxidations<sup>7</sup> is well known in acid media<sup>8</sup>.

**Fig. 4 Plot of  $\log k_1$  vs  $\log [\text{H}^+]$**   
 $[\text{QFC}] = 2.0 \times 10^{-3} \text{ mol dm}^{-3}$ ;  $[\text{unsaturated acid}] = 2.0 \times 10^{-2} \text{ mol dm}^{-3}$ ;  $[\text{NaClO}_4] = 2.0 \times 10^{-1} \text{ mol dm}^{-3}$ ; Solvent : AcOH-H<sub>2</sub>O (50-50% v/v); Temperature : 303 K.



E: ISSN No. 2349-9443

**Effect of Ionic Strength**

Variation of ionic strength of the medium by the addition of NaClO<sub>4</sub> has no pronounced effect on the rate. Similar observations were also reported in the oxidation of unsaturated substrates by QCC<sup>9, 10</sup>.

**Induced Polymerisation**

The reaction mixture failed to initiate polymerisation of acrylonitrile, indicating the absence of formation of free radical species in the reaction sequence.

**Effect of Solvent Composition**

The oxidation of maleic acid and acrylic acid were studied in the solvent containing different proportions of acetic acid and water. The reaction rate increases with increase in acetic acid content of the reaction mixture. The plots of log k<sub>1</sub> against 1/D inverse of dislactric constant were linear with positive

# Asian Resonance

slope. This implies the occurrence of an interaction of an ion-dipole type<sup>11</sup> which confirms the involvement of protonated oxidant species in the rate determining step.

**Effect of Temperature**


The rate increases with the rise in temperature. The reaction between unsaturated acids and QFC were studied at five different temperatures in the range 293K-313K and the activation and thermodynamic parameters were computed by standard procedure (Table 2).The Arrhenius plots of log k<sub>1</sub> versus 1/T were linear (Fig. 5).The entropy of activation are largely negative as expected for bimolecular reaction. The constancy of free energy of activation for both substrates may signal the operation of similar mechanism.

**Table 2. Activation and thermodynamic parameters**

Unsaturated acid	Ea (kJ mol <sup>-1</sup> )	ΔH# (kJ mol <sup>-1</sup> )	ΔG# (kJ mol <sup>-1</sup> )	-ΔS# (JK <sup>-1</sup> mol <sup>-1</sup> )
Maleic acid	55.55	53.03	98.61	150.44
Acrylic acid	55.53	53.01	100.64	157.18

**Fig. 5. Plot of log k<sub>1</sub> vs 1/T**


[QFC]= 2.0×10<sup>-3</sup> mol dm<sup>-3</sup>; [unsaturated acid] = 2.0×10<sup>-2</sup> mol dm<sup>-3</sup>; [HClO<sub>4</sub>] = 0.92 mol dm<sup>-3</sup>; [NaClO<sub>4</sub>] = 2.0×10<sup>-1</sup> mol dm<sup>-3</sup>; Solvent : AcOH-H<sub>2</sub>O (50-50% v/v)



**Mechanism**


It was found that epoxide is the only product of epoxidation of unsaturated acid with QFC.To

explain the formation of product and other observed data, the mechanism in scheme I is proposed.

**Scheme I**

The rate-determining step may be the loose complex formation between the protonated QFC and substrate. An electrophilic attack of Cr(VI), being positively charged in the protonated QFC on the double bond of the substrate produce a four-membered transition state which can yield the


epoxide in the fast step. In the proposed mechanism (Scheme I), a direct chromium to carbon bond does not account for the insensitivity to steric effects very often observed in the oxidation of olefins by Cr(VI)<sup>12</sup>. Thus the most favourable path may be the three centre type addition as depicted in scheme II.

**Scheme II**

Where, R=COOH for maleic acid and R=H for acrylic acid

Both scheme envisage an oxygen atom transfer from the oxidant. This is in accord with the earlier observation made for unsaturated substrates<sup>13, 14</sup>. A closer look on the energy of activation of the reactions under investigation reveals that the values are very close indicating that the electrical and structural properties of solvent system are little affected by the presence of different substrates. The energy of activation of both acids are nearly equal. This shows that both the acids are oxidized with

similar rate. Consequently the rate law is described as follows:



E: ISSN No. 2349-9443

$$\text{Rate} = -\frac{d[\text{QFC}]}{dt} = k_1 [\text{QFCH}^+] [\text{S}]$$

By equation (1)

$$-\frac{d[\text{QFC}]}{dt} = k_1 K [\text{QFC}] [\text{H}^+] [\text{S}]$$

$$-\frac{d[\text{QFC}]}{dt} = k_{\text{obs}} [\text{QFC}] [\text{H}^+] [\text{S}]$$

Where,  $k_1 K = k_{\text{obs}}$ 

The rate law is in accordance with the experimental kinetic findings.

**Endnotes**

1. Cainelli, G. and Cardillo, G. "Chromium Oxidation in Organic Chemistry", Springer-Verlag, New York (1984).
2. Mahanti, M.K. and Banerji, K.K. "Synthetic and Mechanistic aspects of reactions of complexed chromium(VI) compounds", J. Indian Chem. Soc. Vol. 79, pp. 31-44, (2002).
3. Patel, Sabita and Mishra, B.K. "Chromium(VI) oxidants having quaternary ammonium ions: Studies on synthetic applications and oxidation kinetics", Tetrahedron vol. 63, pp. 4367-4406, (2007).
4. Singh, J.V., Awasthi, A., Dipti, Tomar, A. and Singh, D., "Pyridinium and Quinolinium Halochromate : Kinetic and Mechanistic Aspects", Asian J. Chemistry vol. 23, pp. 4744-4750 (2011).
5. Murugesan, V. and Pandurangan, A. "Quinolinium fluorochromate-A New Reagent for the Oxidation of Organic Compounds", Ind. J. Chem., 31B, pp. 377-378, (1992).
6. Pasto, D.J., and Johnson, C.R. "Organic Structure Determination, Prentice-Hall, New Jersey P. 376, (1969).
7. Wiberg, K.B., "Oxidation in Organic Chemistry Part A", Academic Press New York P. 69, (1965).
8. Banerji, K.K. "Kinetics and Mechanism of the oxidation of substituted benzyl alcohol pyridinium chlorochromate" J. Chem. Soc. Res (M), pp. 2561-2564, (1978).
9. Mishra, K., Singh, J.V. and Pandey, A., (6) comparative study on the kinetics of oxidation of unsaturated acids by quinolinium chlorochromate", Proceedings of the National Academy of Sciences, India, vol. 74, pp. 369-378, (2004).
10. Mishra, K., Singh, J.V. and Pandey, A. "Relative Reactivities of some unsaturated compounds towards quinolinium chlorochromate in acetic acid medium in presence of perchloric acid. A kinetic Approach" Oxid. Commun., vol. 27, pp. 90-97, (2004).
11. Amis, E.S., "Solvent Effects on Reaction Rates and Mechanism", Academic Press. New York Press, P. 42 (1967).
12. Rocek, J. and Drozd, J.C. "Evidence for Epoxides as Intermediate in Chromic acid Oxidation of Olefins" J. Am. Chem. Soc., vol. 92, pp. 6668-6669, (1970).
13. Agawal, G.L., and Jain, R., "Kinetics and Mechanism of the Epoxidation of Allyl alcohol with quinolinium fluorochromate" Oxid. Commun., vol. 20, pp. 273-277, (1997).
14. Agarwal, G.L., and Jain, R. "Epoxidation of Crotonaldehyde by Quinolinium Fluorochromate A kinetic study" Oxid. Commun., vol. 22, pp. 514-518, (1999).

# Asian Resonance